

**Table 13. AIDS cases in adolescents and adults under age 25, by sex and exposure category, reported through December 1998, United States**

Male exposure category	13-19 years old				20-24 years old			
	1998		Cumulative total		1998		Cumulative total	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Men who have sex with men	47	(32)	704	(34)	488	(54)	11,079	(62)
Injecting drug use	11	(7)	130	(6)	90	(10)	2,199	(12)
Men who have sex with men and inject drugs	5	(3)	103	(5)	41	(5)	1,853	(10)
Hemophilia/coagulation disorder	16	(11)	740	(36)	23	(3)	623	(4)
Heterosexual contact:	7	(5)	77	(4)	83	(9)	792	(4)
<i>Sex with injecting drug user</i>	—		21		14		256	
<i>Sex with person with hemophilia</i>	—		1		—		4	
<i>Sex with transfusion recipient with HIV infection</i>	—		—		—		10	
<i>Sex with HIV-infected person, risk not specified</i>	7		55		69		522	
Receipt of blood transfusion, blood components, or tissue	8	(5)	90	(4)	2	(0)	105	(1)
Risk not reported or identified <sup>1</sup>	53	(36)	231	(11)	180	(20)	1,146	(6)
Male subtotal	147	(100)	2,075	(100)	907	(100)	17,797	(100)
<b>Female exposure category</b>								
Injecting drug use	14	(9)	190	(14)	87	(15)	1,827	(28)
Hemophilia/coagulation disorder	1	(1)	12	(1)	—	—	13	(0)
Heterosexual contact:	61	(41)	708	(53)	289	(49)	3,614	(54)
<i>Sex with injecting drug user</i>	13		255		62		1,463	
<i>Sex with bisexual male</i>	3		37		11		255	
<i>Sex with person with hemophilia</i>	—		13		3		51	
<i>Sex with transfusion recipient with HIV infection</i>	—		2		1		22	
<i>Sex with HIV-infected person, risk not specified</i>	45		401		212		1,823	
Receipt of blood transfusion, blood components, or tissue	3	(2)	88	(7)	2	(0)	114	(2)
Risk not reported or identified	71	(47)	350	(26)	216	(36)	1,072	(16)
Female subtotal	150	(100)	1,348	(100)	594	(100)	6,640	(100)
<b>Total</b>	<b>297</b>		<b>3,423</b>		<b>1,501</b>		<b>24,437</b>	

<sup>1</sup>See figure 6.